

AG
T

*Algebraic & Geometric
Topology*

Volume 25 (2025)

**Coarse and bi-Lipschitz embeddability of subspaces
of the Gromov–Hausdorff space into Hilbert spaces**

NICOLÒ ZAVA

Coarse and bi-Lipschitz embeddability of subspaces of the Gromov–Hausdorff space into Hilbert spaces

NICOLÒ ZAVA

We discuss the embeddability of subspaces of the Gromov–Hausdorff space, which consists of isometry classes of compact metric spaces endowed with the Gromov–Hausdorff distance, into Hilbert spaces. These embeddings are particularly valuable for applications to topological data analysis. We prove that its subspace consisting of metric spaces with at most n points has asymptotic dimension $n(n - 1)/2$. Thus, there exists a coarse embedding of that space into a Hilbert space. On the contrary, if the number of points is not bounded, then the subspace cannot be coarsely embedded into any uniformly convex Banach space and so, in particular, into any Hilbert space. Furthermore, we prove that, even if we restrict to finite metric spaces whose diameter is bounded by some constant, the subspace still cannot be bi-Lipschitz embedded into any finite-dimensional Hilbert space. We obtain both nonembeddability results by finding obstructions to coarse and bi-Lipschitz embeddings in families of isometry classes of finite subsets of the real line endowed with the Euclidean–Hausdorff distance.

46B85, 51F30; 54B20

1 Introduction

The Gromov–Hausdorff distance d_{GH} measures how two metric spaces resemble each other. It was introduced by Edwards in [27], and then rediscovered and generalised by Gromov [31]. Until around 2000, the Gromov–Hausdorff distance had been mainly used by pure mathematicians who were interested in the induced topology. That direction is still of great interest, and, as an example, we mention the two recent papers [3; 4].

In addition to the intrinsic interest in it, a great impulse to study the quantitative aspects of the Gromov–Hausdorff distance came from its applications in topological data analysis, which is a fast-growing subject aiming to use topological techniques to analyse a wide range of real-world data (see, for example, [13; 33], and [30] for a growing dataset of real-world applications). The Gromov–Hausdorff distance provides a theoretical framework to directly compare point clouds by considering them as metric spaces. This approach proved to be useful in shape recognition and comparison [43; 46; 47], which arises, for example, in molecular biology, databases of objects, face recognition and matching of articulated objects.

Comparing two metric spaces using the Gromov–Hausdorff distance directly is computationally expensive. Even approximating it within a factor of 3 for trees with unit edge length is NP-hard ([2; 59]; see also [43],

where the author discussed the connection between computing the Gromov–Hausdorff distance and a class of NP-hard problems). Therefore, creating efficiently computable invariants to approximate the Gromov–Hausdorff distance is of particular interest. Following [45], an *invariant* ψ associates to a metric space X an element $\psi(X)$ of another metric space $(\mathfrak{Y}, d_{\mathfrak{Y}})$ in such a way that, if X and Y are two isometric metric spaces, then $\psi(X) = \psi(Y)$. Furthermore, an invariant ψ is *stable* if there exists a function $\rho_+ : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ such that

$$(1) \quad d_{\mathfrak{Y}}(\psi(X), \psi(Y)) \leq \rho_+(d_{\text{GH}}(X, Y))$$

for every pair of metric spaces X and Y . Stability implies that small perturbations of the metric spaces have a limited effect on the associated invariants. Therefore, considering similarity recognition, we avoid false negatives, which are situations where two metric spaces are very similar in the Gromov–Hausdorff distance, but their invariants are far apart. Furthermore, stable invariants can be used to provide lower bounds to the Gromov–Hausdorff distance as shown in [45]. We refer to the latter paper for a wide range of stable invariants. Additional examples are hierarchical clustering [14; 15] and persistence diagrams induced by the Vietoris–Rips, the Dowker, and the Čech filtrations ([17; 18]; see also [25] for details and applications of persistent homology).

In contrast to false negatives, even though still undesirable, it is often acceptable when an invariant produces false positives, where two dissimilar spaces are mapped to close values. In this paper, we study when stable invariants are actually bound to lose information because of the unavoidable creation of false positives. We focus our study on those invariants taking values in a Hilbert space. Those are particularly relevant for the applications in machine learning pipelines since many algorithms expect either data in the form of Euclidean vectors or at least access to a so-called feature map into a Hilbert space. As formally stated in the sequel, we prove that the existence of a stable invariant avoiding false positives strongly depends on a bound on the cardinality of the metric spaces.

Our approach to the problem requires notions and techniques from coarse geometry. Intuitively, this field, also known as large-scale geometry, focuses on large-scale, global properties of spaces ignoring local features. We refer to [52; 58] for a wide introduction. A map $\psi : (X, d_X) \rightarrow (Y, d_Y)$ between two metric spaces is said to be a *coarse embedding* if there exist two maps $\rho_-, \rho_+ : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ such that $\rho_- \rightarrow \infty$ and, for every $x, y \in X$,

$$(2) \quad \rho_-(d_X(x, y)) \leq d_Y(\psi(x), \psi(y)) \leq \rho_+(d_X(x, y)).$$

In the case of stable invariants, ie, satisfying (1), a lower bound as in (2) prevents false positives since the larger the Gromov–Hausdorff distance, the larger the distance between the two associated invariants.

Coarse embeddings have been introduced by Gromov and extensively studied in coarse geometry. A crucial application of this theory is due to Yu, who proved in [65] that those metric spaces that can be coarsely embedded into a Hilbert space satisfy the Novikov and the coarse Baum–Connes conjectures generalising results contained in [64]. This result motivated two research directions. On one hand, since

an explicit coarse embedding can be hard to construct, a plethora of conditions ensuring its existence have been defined and investigated. We refer the interested reader to [52] for a discussion on the topic and to [63] for more examples. Among these properties, if a space has finite asymptotic dimension, then it can be coarsely embedded into a Hilbert space [34; 65]. Asymptotic dimension is a large-scale counterpart of Lebesgue’s covering dimension introduced in [32] (see also [7]). On the other hand, examples of metric spaces that cannot be coarsely embedded were constructed, for example, in [22; 39]. Showing that one of those pathological examples can be coarsely embedded into a metric space X is a technique to prove that X itself cannot be coarsely embedded into any Hilbert space.

Those two strategies have been adopted to prove if metric spaces emerging in different fields can be coarsely embedded into Hilbert spaces. In topological data analysis, collections of persistence diagrams endowed with various metrics represent a prominent example. It was proved in [48] that the space of persistence diagrams of at most n points endowed with the Hausdorff distance has finite asymptotic dimension, and so it can be coarsely embedded into a Hilbert space. This result, despite being nonconstructive, motivated further research in that direction that eventually led to explicit bi-Lipschitz and coarse embeddings in [6] and [49], respectively. In the opposite direction, it was proved in [11; 48; 62] that spaces of persistence diagrams with various metrics cannot be coarsely embedded into any Hilbert space. We also refer to [55], where the authors showed the equivalence of this problem with the embeddability of Wasserstein spaces.

Another example can be found in [29]. Motivated by the interest in invariants in crystallography and pharmaceuticals (see [26]), the authors used the previously described strategy to prove that spaces of periodic point sets equipped with the Euclidean bottleneck distance cannot be coarsely embedded into any uniformly convex Banach space. The constructions used in that paper are adapted from those developed in [63] to prove the analogous noncoarse embeddability results for families of finite subsets of metric spaces endowed with the Hausdorff distance.

In this paper, we prove the following results.

Theorem A *The space $\mathcal{GH}^{\leq n}$ of isometry classes of metric spaces with at most n points endowed with the Gromov–Hausdorff distance has asymptotic dimension $n(n-1)/2$, and so it can be coarsely embedded into a Hilbert space.*

Theorem B *The space $\mathcal{GH}^{<\omega}$ of isometry classes of finite metric spaces endowed with the Gromov–Hausdorff distance cannot be coarsely embedded into any uniformly convex Banach space, and so, in particular, into any Hilbert space.*

As an immediate consequence of **Theorem B**, the same result holds for the *Gromov–Hausdorff space* \mathcal{GH} , which is the metric space of isometry classes of compact metric spaces equipped with the Gromov–Hausdorff distance. In the paper, we prove a stronger version of **Theorem B** stating that already the much smaller subspace of $\mathcal{GH}^{<\omega}$ consisting of all isometry classes of finite subsets of the real line cannot be

coarsely embedded into any uniformly convex Banach space (Theorem 4.2). Thanks to a recent result due to Majhi, Vitter and Wenk [42], Theorem 4.2, and therefore also Theorem B, will follow from the fact that an obstruction to coarse embeddability is found in the space of isometry classes of finite subsets of the real line endowed with the Euclidean–Hausdorff distance (a modification of the Gromov–Hausdorff distance for subsets of \mathbb{R}^d).

We conclude the paper focussing on the subspace $\mathcal{GH}_{\leq R}^{<\omega}$ of $\mathcal{GH}^{<\omega}$ whose elements have diameter bounded by a constant $R > 0$. Since the diameter of this subspace is bounded, the map collapsing the space into a point is trivially a coarse embedding. An immediate follow-up question is whether it can be bi-Lipschitz embedded, as in the case of persistence diagrams with at most n points. We provide a partial negative answer.

Theorem C $\mathcal{GH}_{\leq R}^{<\omega}$ *cannot be bi-Lipschitz embedded into any finite-dimensional Hilbert space.*

Inspired by [16], where the authors proved that certain spaces of persistence diagrams cannot be bi-Lipschitz embedded into any finite-dimensional Hilbert space, we compute the Assouad dimension ([5]; see also [10] for an earlier definition) of $\mathcal{GH}_{\leq R}^{<\omega}$ and show that it is infinite. This dimension notion was in fact introduced to provide such embeddability obstructions. More precisely, we show that already the subset consisting of all isometry classes of finite subsets of an interval has infinite Assouad dimension and cannot be bi-Lipschitz embedded into any finite-dimensional Hilbert space. Again, using the aforementioned Majhi, Vitter and Wenk’s theorem, we deduce our claims from the analogous results for the space of isometry classes of finite subsets of an interval endowed with the Euclidean–Hausdorff distance (Theorem 5.1 and Proposition 5.3).

The paper is organised as follows. In Section 2 we provide the needed background regarding the Gromov– and the Euclidean–Hausdorff distances. In Section 3, the asymptotic dimension is introduced and Theorem A is proved. Theorem B is shown in Section 4, and, finally, we define the Assouad dimension and provide Theorem C in Section 5. We conclude the paper discussing a list of questions in Section 5.1.

Acknowledgements The author was supported by the FWF Grant, Project number I4245-N35. The author would like to thank Thomas Weighill for the helpful discussions around Theorem 3.10, and Takamitsu Yamauchi for bringing to my attention the fundamental reference [35]. Furthermore, the author is thankful for the detailed and helpful comments of the reviewer of this manuscript.

Notation We denote by \mathbb{N} , \mathbb{Q} and \mathbb{R} the set of natural numbers including 0, the set of rational numbers, and the set of real numbers, respectively. For $c \in \mathbb{R}$, we also write

$$\mathbb{R}_{\geq c} = \{x \in \mathbb{R} \mid x \geq c\} \quad \text{and} \quad \mathbb{R}_{>c} = \{x \in \mathbb{R} \mid x > c\}.$$

For a set X , we denote by $|X|$ its cardinality. Moreover, for $n \in \mathbb{N}$, we define the following subsets of the power set of X :

$$[X]^{=n} = \{A \subseteq X \mid |A| = n\}, \quad [X]^{\leq n} = \bigcup_{k \leq n} [X]^{=n}, \quad [X]^{<\omega} = \bigcup_{k \in \mathbb{N}} [X]^{=n}.$$

2 The Gromov–Hausdorff distance and the Euclidean–Hausdorff distance

We recall some basic notions, the definitions of the Gromov–Hausdorff and the Euclidean–Hausdorff distances and their relationships. We refer to [12; 44; 45; 54; 61] for comprehensive discussions on the Gromov–Hausdorff distance.

Definition 2.1 A pair (X, d) consisting of a set X and a map $d: X \times X \rightarrow \mathbb{R}$ is called a *network* [19]. A network (X, d) is a *metric space* (and d is a *metric*) if it satisfies

- (M1) for every $x, y \in X$, $d(x, y) \geq 0$ and $d(x, x) = 0$;
- (M2) for every $x, y \in X$, $d(x, y) = 0$ if and only if $x = y$;
- (M3) for every $x, y \in X$, $d(x, y) = d(y, x)$;
- (M4) for every $x, y, z \in X$, $d(x, y) \leq d(x, z) + d(z, y)$.

Let us recall that an *isometry* between two networks (X, d_X) and (Y, d_Y) is a bijective map $\psi: X \rightarrow Y$ such that, for every $x, x' \in X$, $d_Y(\psi(x), \psi(x')) = d_X(x, x')$. In that case, X and Y are said to be *isometric*.

For a subset A of a metric space (X, d) , its *diameter* is

$$\text{diam } A = \sup_{x, y \in A} d(x, y).$$

A *correspondence* \mathcal{R} between two sets X and Y is a relation $\mathcal{R} \subseteq X \times Y$ such that every $x \in X$ is in relation with at least one element $y \in Y$ and vice versa. Then, for every metric space (Z, d) , the *Hausdorff distance* is defined as follows: for every $X, Y \subseteq Z$,

$$d_H(X, Y) = \inf_{\mathcal{R} \subseteq X \times Y \text{ correspondence}} \sup_{(x, y) \in \mathcal{R}} d(x, y).$$

Definition 2.2 Given two metric spaces X and Y , their *Gromov–Hausdorff distance* d_{GH} is

$$d_{\text{GH}}(X, Y) = \inf_{Z \text{ metric space}} \inf \{d_H(i_X(X), i_Y(Y)) \mid i_X: X \rightarrow Z \text{ and } i_Y: Y \rightarrow Z \text{ isometric embeddings}\}.$$

The reader may notice an abuse of notation in the previous definition since all possible metric spaces form a proper class. However, the infimum value can be achieved by investigating just a set of spaces. Indeed, it is enough to consider the disjoint union $X \sqcup Y$ endowed with pseudometrics (where the distance between distinct points may be zero) whose restrictions to the subsets X and Y coincide with the original metrics. We refer to [12] for the details.

If two metric spaces are isometric, their Gromov–Hausdorff distance is 0. The converse implication does not hold in general. However, if X and Y are compact and $d_{\text{GH}}(X, Y) = 0$, then X and Y are isometric.

Denote by \mathcal{GH} the set of all isometry classes of compact metric spaces endowed with d_{GH} , where the Gromov–Hausdorff distance between two isometry classes is the Gromov–Hausdorff distance between

any pair of representatives. Since two compact metric spaces are isometric if and only if their Gromov–Hausdorff distance is 0 (see, for example, [12]), \mathcal{GH} is a metric space, also called the *Gromov–Hausdorff space*. Furthermore, we consider the subspace $\mathcal{GH}^{<\omega}$ of \mathcal{GH} consisting of isometry classes of finite metric space, which is dense in \mathcal{GH} . Actually, the subspace $\mathcal{GH}_{\mathbb{Q}}^{<\omega}$ of isometry classes of finite spaces endowed with metrics taking values in \mathbb{Q} is dense in \mathcal{GH} [54]. Therefore, \mathcal{GH} is separable.

The Gromov–Hausdorff distance can be alternatively characterised using correspondences. If (X, d_X) and (Y, d_Y) are two networks, and $\mathcal{R} \subseteq X \times Y$ is a correspondence between them, the *distortion* of \mathcal{R} is

$$\text{dis } \mathcal{R} = \sup_{(x_1, y_1), (x_2, y_2) \in \mathcal{R}} |d_X(x_1, x_2) - d_Y(y_1, y_2)|.$$

Definition 2.3 [19] Let X and Y be two networks. Then their *network distance* is

$$d_{\mathcal{N}}(X, Y) = \frac{1}{2} \inf_{\mathcal{R} \subseteq X \times Y \text{ correspondence}} \text{dis } \mathcal{R}.$$

It is known that, if X and Y are metric spaces, then $d_{\mathcal{GH}}(X, Y) = d_{\mathcal{N}}(X, Y)$ (see, for example, [12]). A further characterisation of the Gromov–Hausdorff distance can be found in [36].

In [66], a characterisation of the network distance for *quasimetric spaces* (ie, networks satisfying (M1), (M2) and (M4)) in the spirit of [Definition 2.2](#) is provided.

The Gromov–Hausdorff distance is difficult to compute even in simple cases. For example, the distance between spheres of different dimensions endowed with their geodesic distance is not known in general [1; 41]. To approximate it, it is convenient to consider another related distance.

Definition 2.4 Let X and Y be two subsets of \mathbb{R}^d . Consider them as metric spaces. Then, their *Euclidean–Hausdorff distance* d_{EH} is defined as

$$d_{\text{EH}}(X, Y) = \inf \{d_{\text{H}}(i_X(X), i_Y(Y)) \mid i_X: X \rightarrow \mathbb{R}^d \text{ and } i_Y: Y \rightarrow \mathbb{R}^d \text{ isometric embeddings}\}.$$

Using the following folklore result (see, for example, [9, Chapter IV, Section 38]), d_{EH} can be conveniently characterised.

Theorem 2.5 If $f: X \rightarrow Y$ is an isometry between two subsets of \mathbb{R}^d , then there exists an isometry $\tilde{f}: \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that $\tilde{f}|_X = f$.

Corollary 2.6 (see, for example, [3, Corollary 4.3]) If X and Y are two subsets of \mathbb{R}^d , then

$$d_{\text{EH}}(X, Y) = \inf_{f \in \text{Isom}(\mathbb{R}^d)} d_{\text{H}}(X, f(Y)),$$

where $\text{Isom}(\mathbb{R}^d)$ denotes the group of isometries of \mathbb{R}^d .

Clearly, $d_{\text{EH}}(X, Y) \geq d_{\text{GH}}(X, Y)$ for every pair X and Y of subsets of an Euclidean space \mathbb{R}^d . Moreover, the inequality can be strict (for example, see [44]). A lower bound on the Gromov–Hausdorff distance depending on the Euclidean–Hausdorff distance was provided in [44].

Theorem 2.7 *For every pair of compact subsets X and Y of \mathbb{R}^d ,*

$$d_{\text{GH}}(X, Y) \leq d_{\text{EH}}(X, Y) \leq c_d \sqrt{M \cdot d_{\text{GH}}(X, Y)},$$

where $M = \max\{\text{diam } X, \text{diam } Y\}$ and c_d is a constant depending only on the dimension d .

However, if X and Y are finite subsets of \mathbb{R} , linear lower bounds to d_{GH} depending on d_{EH} can be proved.

Theorem 2.8 [42, Theorem 3.2] *For every pair X and Y of compact subsets of \mathbb{R} ,*

$$\frac{4}{5}d_{\text{EH}}(X, Y) \leq d_{\text{GH}}(X, Y) \leq d_{\text{EH}}(X, Y).$$

If we denote by \mathcal{EH}_1 ($\mathcal{EH}_1^{<\omega}$) the space of isometry classes of compact (finite, respectively) subsets of the real line endowed with the Euclidean–Hausdorff distance, the canonical inclusion of \mathcal{EH}_1 into \mathcal{GH} and that of $\mathcal{EH}_1^{<\omega}$ into $\mathcal{GH}^{<\omega}$ are bi-Lipschitz according to Theorem 2.8. Let us recall that a map $\psi: X \rightarrow Y$ between metric spaces is a *bi-Lipschitz embedding* if there are two linear maps $\rho_-: x \mapsto a \cdot x$ and $\rho_+: x \mapsto b \cdot x$, where $a, b > 0$, satisfying (2).

3 The space of metric spaces of at most n points is coarsely embeddable into a Hilbert space

Given two subsets Y, Z of a metric space (X, d) and a radius $r \geq 0$ we write

$$\text{dist}_d(Y, Z) = \text{dist}(Y, Z) = \inf\{d(y, z) \mid y \in Y, z \in Z\} \quad \text{and} \quad B_d(Y, r) = \bigcup_{y \in Y} B_d(y, r),$$

where $B_d(y, r)$ denotes the closed ball centred in y with radius r .

A family of subsets \mathcal{U} of a metric space X is said to be

- *uniformly bounded* if there exists $R \geq 0$ such that $\text{diam } U \leq R$ for every $U \in \mathcal{U}$ (if we need to specify R , we say it is *R-bounded*);
- *r-disjoint* for some $r > 0$ if $\text{dist}(U, V) > r$ for every $U, V \in \mathcal{U}$ with $U \neq V$.

Definition 3.1 [32] Let X be a metric space. The *asymptotic dimension* of X is at most $n \in \mathbb{N}$ (and we write $\text{asdim } X \leq n$) if for every $r > 0$ there exists a uniformly bounded cover $\mathcal{U} = \mathcal{U}_0 \cup \dots \cup \mathcal{U}_n$ of X such that \mathcal{U}_i is r -disjoint for every $i = 0, \dots, n$. Furthermore, $\text{asdim } X = n$ if $\text{asdim } X \leq n$ and $\text{asdim } X \not\leq n$, and $\text{asdim } X = \infty$ if $\text{asdim } X \not\leq m$ for every $m \in \mathbb{N}$.

Example 3.2 (see [52]) For every $n \in \mathbb{N}$, $\text{asdim } \mathbb{R}^n = \text{asdim}(\mathbb{R}_{\geq 0})^n = n$ where \mathbb{R}^n is equipped with any p -norm, $p \in [1, \infty]$, and $(\mathbb{R}_{\geq 0})^n$ with any of the inherited metrics.

Let us recall two basic properties of the asymptotic dimension that we are going to use later in this section.

Proposition 3.3 (see [7]) *Let (X, d) be a metric space and $Y \subseteq X$ be a subspace. Then*

- (a) $\text{asdim } Y \leq \text{asdim } X$, and
- (b) $\text{asdim } Y = \text{asdim } X$ provided that Y is **large** in X , ie, there exists $r \geq 0$ such that $B_d(Y, r) = X$.

The goal of this section is to prove [Theorem A](#), which we obtain as a particular case of the more general [Corollary 3.11](#).

Lemma 3.4 *For every $n \in \mathbb{N}$, $\text{asdim } \mathcal{GH}^{\leq n} \geq n(n-1)/2$.*

Proof By [35, Theorem 4.1], $\mathcal{GH}^{\leq n}$ contains isometric copies of arbitrarily large balls of $\mathbb{R}^{n(n-1)/2}$ endowed with the supremum metric. Then, [48, Lemma 2.10] implies that $\text{asdim } \mathcal{GH}^{\leq n} \geq n(n-1)/2$. \square

In order to prove the opposite inequality, we need to show different steps. Let us start with recalling a known result.

Theorem 3.5 ([56]; see also [37, Theorem 4.6]) *For every $n \in \mathbb{N}$, $\text{asdim}([X]^{\leq n}, d_H) \leq n \text{ asdim } X$. In particular, $\text{asdim}[\mathbb{R}]^{\leq n} \leq n$ and $\text{asdim}[\mathbb{R}_{\geq 0}]^{\leq n} \leq n$.*

For every positive integer $n \in \mathbb{N}$, let us define \mathcal{X}_n as the metric subspace of $([\mathbb{R}_{\geq 0}]^{\leq n+1}, d_H)$ whose elements contain the point 0. According to [Theorem 3.5](#), $\text{asdim } \mathcal{X}_n \leq n+1$ since the asymptotic dimension is monotone ([Proposition 3.3\(a\)](#)). In the sequel, we improve that bound showing that $\text{asdim } \mathcal{X}_n \leq n$. The proof outline is similar to and inspired by that of [48, Theorem 3.2]. First, we need a classical preliminary result.

Given two families \mathcal{U} and \mathcal{V} of subsets of a metric space X and $r > 0$, we define a new family of subsets as

$$\mathcal{U} \cup_r \mathcal{V} = \{N_r(U, V) \mid U \in \mathcal{U}\} \cup \{V \in \mathcal{V} \mid \forall U \in \mathcal{U}, \text{dist}(V, U) > r\},$$

where $N_r(U, \mathcal{V}) = U \cup \bigcup \{V \in \mathcal{V} \mid \text{dist}(U, V) \leq r\}$.

Let us immediately note that $\bigcup (\mathcal{V} \cup_r \mathcal{U}) \supseteq \bigcup \mathcal{V} \cup \bigcup \mathcal{U}$.

Lemma 3.6 [7, Proposition 24] *Let \mathcal{U} be an r -disjoint, R -bounded family of subsets of X with $R \geq r$. Let \mathcal{V} be a $5R$ -disjoint, uniformly bounded family of subsets of X . Then $\mathcal{V} \cup_r \mathcal{U}$ is r -disjoint and uniformly bounded.*

Lemma 3.7 *For every $n \in \mathbb{N}$, $\text{asdim } \mathcal{X}_n \leq n$.*

Proof Let us prove the result by induction.

If $n = 1$, then, for every $r > 0$, the usual uniformly bounded cover used to prove that the real line has asymptotic dimension 1 (see [7; 52]) can be adapted. More precisely, for every $r > 0$, define, for $i = 0, 1$,

$$\mathcal{U}_i = \{\{0, x\} \mid x \in V_k^i\} \mid k \in \mathbb{N}\}, \quad \text{where } V_k^i = [(4k+2i)r, (4k+2i+2)r] \subseteq \mathbb{R}_{\geq 0}$$




Figure 1: A representation of the uniformly bounded cover $\mathcal{V} = \mathcal{V}_0 \cup \mathcal{V}_1$ showing that $\text{asdim } \mathbb{R}_{\geq 0} \leq 1$. In red, the elements of $\mathcal{V}_0 = \{V_k^0 \mid k \in \mathbb{N}\}$ and, in blue, the subsets contained in the family $\mathcal{V}_1 = \{V_k^1 \mid k \in \mathbb{N}\}$.

(see Figure 1). Then, each \mathcal{U}_i is r -disjoint and $\mathcal{U}_0 \cup \mathcal{U}_1$ forms a uniformly bounded cover of \mathcal{X}_1 . Hence, $\text{asdim } \mathcal{X}_1 \leq 1$.

Suppose now that the assertion is true for some $n \in \mathbb{N}$. We want to show that $\text{asdim } \mathcal{X}_{n+1} \leq n+1$. Fix a positive $r > 0$. Then we can write

$$\mathcal{X}_{n+1} = \widetilde{\mathcal{X}_{n+1}} \cup B_{d_H}(\mathcal{X}_n, r),$$

where $\widetilde{\mathcal{X}_{n+1}}$ consists of subsets A of $\mathbb{R}_{\geq 0}$ such that, if $x \in A$ has $|x| \leq r$, then $x = 0$. It indeed represents a partition. In fact, every element $C \in \mathcal{X}_{n+1}$ can be decomposed $C = C_0 \cup C_1$ where $\max_{x \in C_0} |x| \leq r$ and $\min_{x \in C_1} |x| > r$. Then, either $C_0 = \{0\}$ and so $C \in \widetilde{\mathcal{X}_{n+1}}$, or $d_H(C, C_1 \cup \{0\}) \leq r$ and $C_1 \cup \{0\} \in \mathcal{X}_n$.

Let us now consider $\widetilde{\mathcal{X}_{n+1}}$, and define the subspace $\mathcal{Y}_n = \{X \setminus \{0\} \mid X \in \widetilde{\mathcal{X}_{n+1}}\}$ of $[\mathbb{R}_{\geq 0}]^{\leq n+1}$. Monotonicity of the asymptotic dimension (Proposition 3.3(a)) implies that $\text{asdim } \mathcal{Y}_n \leq n+1$. Hence, there exists a uniformly bounded cover $\mathcal{W} = \mathcal{W}_0 \cup \dots \cup \mathcal{W}_{n+1}$ of \mathcal{Y}_n such that \mathcal{W}_i is r -disjoint for every $i = 0, \dots, n+1$. Let $R \geq r$ be an upper bound to the diameter of the elements in \mathcal{W} . For every $i = 0, \dots, n+1$, construct the family of subsets $\widetilde{\mathcal{W}_i} = \{W \cup \{0\} \mid W \in \mathcal{W}_i\}$ of $\widetilde{\mathcal{X}_{n+1}}$. Note that $\widetilde{\mathcal{W}} = \widetilde{\mathcal{W}_0} \cup \dots \cup \widetilde{\mathcal{W}_{n+1}}$ is a cover of $\widetilde{\mathcal{X}_{n+1}}$. Furthermore, each element of $\widetilde{\mathcal{W}}$ has diameter bounded by R and it is easy to see that each of the families $\widetilde{\mathcal{W}_i}$, $i = 0, \dots, n+1$, is r -disjoint.

Since \mathcal{X}_n is large in $B_{d_H}(\mathcal{X}_n, r)$, $\text{asdim } B_{d_H}(\mathcal{X}_n, r) \leq n \leq n+1$ by the inductive hypothesis and Proposition 3.3(b). Hence, there exists a uniformly bounded cover $\mathcal{V} = \mathcal{V}_0 \cup \dots \cup \mathcal{V}_{n+1}$ of $B_{d_H}(\mathcal{X}_n, r)$ such that \mathcal{V}_i is $5R$ -disjoint for every $i = 0, \dots, n+1$.

Define, for every $i = 0, \dots, n+1$,

$$\mathcal{U}_i = \widetilde{\mathcal{W}_i} \cup_r \mathcal{V}_i.$$

Hence, $\mathcal{U} = \mathcal{U}_0 \cup \dots \cup \mathcal{U}_{n+1}$ is a uniformly bounded cover of $\mathcal{X}_{n+1} = \bigcup \widetilde{\mathcal{W}} \cup \bigcup \mathcal{V}$ and \mathcal{U}_i is r -disjoint for every $i = 0, \dots, n+1$ according to Lemma 3.6. Thus, $\text{asdim } \mathcal{X}_{n+1} \leq n+1$. \square

On the family of all isometry classes of finite networks, the network distance d_N is a *pseudometric*, ie, it satisfies the properties (M1), (M3) and (M4) stated in Definition 2.1 [19]. Consider the equivalence relation between finite networks given by $X \sim Y$ if $d_N(X, Y) = 0$. We refer to [20] where this equivalence relation is completely characterised. The space obtained by quotienting the family of all isometry classes of finite networks under this equivalence relation is a metric space (it is a standard way to obtain a metric

space out of a pseudometric space; see, for example, [12, Proposition 1.1.5]). We denote it by $\mathcal{N}^{<\omega}$. Let us recall that the distance between two elements in $\mathcal{N}^{<\omega}$ is the network distance between any two representatives. For the sake of simplicity, we will be directly working with representatives of objects in $\mathcal{N}^{<\omega}$ instead of their equivalence classes without explicit mention. Let us also consider the following metric subspaces of $\mathcal{N}^{<\omega}$:

- $\mathcal{S}^{<\omega}$ – the family of equivalence classes of finite *pseudo-semi-metric spaces*, ie, networks satisfying (M1) and (M3);
- $\mathcal{N}^{\leq n}$ – the family of equivalence classes of networks whose cardinality is at most n ;
- $\mathcal{S}^{\leq n} = \mathcal{S}^{<\omega} \cap \mathcal{N}^{\leq n}$.

Define the map $\mathcal{D}: \mathcal{N}^{<\omega} \rightarrow [\mathbb{R}]^{<\omega}$ as follows: for every $(X, d) \in \mathcal{N}^{<\omega}$,

$$\mathcal{D}(X) = d(X \times X) \subseteq \mathbb{R}.$$

The subset $\mathcal{D}(X)$ is called the *distance set* of X . Note that $\mathcal{D}|_{\mathcal{S}^{\leq n}}: \mathcal{S}^{\leq n} \rightarrow \mathcal{X}_{n(n-1)/2}$. For the sake of simplicity, we denote by \mathcal{D} also the restriction.

Proposition 3.8 [20, Proposition 4.3.4] *The map $\mathcal{D}: \mathcal{N}^{<\omega} \rightarrow ([\mathbb{R}]^{<\omega}, d_H)$ is well defined and 2-Lipschitz (ie, $d_H(\mathcal{D}(X), \mathcal{D}(Y)) \leq 2d_{\mathcal{N}}(X, Y)$ for every $X, Y \in \mathcal{N}^{<\omega}$).*

Proof Let $(X, d_X), (Y, d_Y) \in \mathcal{N}^{<\omega}$ and \mathcal{R} be a correspondence such that $R = \text{dis } \mathcal{R} = 2d_{\mathcal{N}}(X, Y)$. For every $x_1, x_2 \in X$ pick $y_1, y_2 \in Y$ satisfying $(x_i, y_i) \in \mathcal{R}$, for $i = 1, 2$. Then, $|d_X(x_1, x_2) - d_Y(y_1, y_2)| \leq R$. Since the same argument can be carried out also for every pair of points $y_1, y_2 \in Y$, $d_H(\mathcal{D}(X), \mathcal{D}(Y)) \leq R$. This inequality also implies that the map is well defined. \square

Even if we restrict ourselves to consider only metric spaces that are finite subsets of the real line, the map \mathcal{D} is not injective [8]. However, it has a further property that allows us to deduce an upper bound to the asymptotic dimension of $\mathcal{GH}^{\leq n}$.

Let us recall that, for a positive integer $k \in \mathbb{N}$, a map $\varphi: (X, d_X) \rightarrow (Y, d_Y)$ between metric spaces is *coarsely k -to-1* [50] if

- there exists a map $\rho_+: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ such that, for every $x, y \in X$,

$$d_Y(\varphi(x), \varphi(y)) \leq \rho_+(d_X(x, y))$$

(ie, φ is *bornologous*);

- for every $R \geq 0$, there exists $S \geq 0$ such that for every $y \in Y$ there are $x_1, \dots, x_k \in X$ satisfying

$$\varphi^{-1}(B_{d_Y}(y, R)) \subseteq \bigcup_{i=1}^k B_{d_X}(x_i, S).$$

Coarsely k -to-1 maps play a very important role in providing bounds for the asymptotic dimension.

Theorem 3.9 ([23; 50]; see also [24]) *Let $\varphi: X \rightarrow Y$ be a coarsely k -to-1 surjective map between metric spaces. Then,*

$$\text{asdim } X \leq \text{asdim } Y \leq k(\text{asdim } X + 1) - 1.$$

Theorem 3.10 *The map $\mathcal{D}: \mathcal{S}^{\leq n} \rightarrow \mathcal{D}(\mathcal{S}^{\leq n}) = \mathcal{X}_{n(n-1)/2}$ is coarsely k -to-1 for some suitable k .*

Proof Proposition 3.8 implies that the map is bornologous.

Let us fix $R \in \mathbb{R}_{\geq 0}$ and $D \in \mathcal{D}(\mathcal{S}^{\leq n}) = \mathcal{X}_{n(n-1)/2}$. In particular, $0 \in D$. Suppose that $(Y, d_Y) \in \mathcal{S}^{\leq n}$ satisfies $d_H(D, \mathcal{D}(Y)) \leq R$. Then, there exists a function $f: \mathcal{D}(Y) \rightarrow D$ such that $|f(a) - a| \leq R$ for every $a \in \mathcal{D}(Y)$. Furthermore, without loss of generality, we can require that $f(0) = 0$ (note that $0 \in \mathcal{D}(Y)$).

Define a new object $X_Y \in \mathcal{S}^{\leq n}$ as $X_Y = (Y, f \circ d_Y)$. Then,

- (a) $X_Y \in \mathcal{S}^{\leq n}$;
- (b) $\mathcal{D}(X_Y) \subseteq D$;
- (c) $d_N(Y, X_Y) \leq \frac{1}{2} \text{dis id} \leq \frac{1}{2}R$.

Items (a) and (b) imply that

$$|\{X_Y \mid Y \in \mathcal{S}^{\leq n} : d_H(\mathcal{D}(Y), D) \leq R\}| \leq |D|^{\frac{1}{2}n(n-1)} \leq \left(\frac{1}{2}n(n-1) + 1\right)^{\frac{1}{2}n(n-1)} =: k.$$

Hence, \mathcal{D} is a coarsely k -to-1 map. □

Corollary 3.11 *For every $n \in \mathbb{N}$, $\text{asdim } \mathcal{GH}^{\leq n} = \text{asdim } \mathcal{S}^{\leq n} = n(n-1)/2$.*

Proof The claim is implied by

$$\frac{1}{2}n(n-1) \leq \text{asdim } \mathcal{GH}^{\leq n} \leq \text{asdim } \mathcal{S}^{\leq n} \leq \text{asdim } \mathcal{D}(\mathcal{S}^{\leq n}) = \text{asdim } \mathcal{X}_{\frac{1}{2}n(n-1)} \leq \frac{1}{2}n(n-1),$$

where the first inequality is stated in Lemma 3.4, the second one follows from Proposition 3.3(a), the third one from Theorem 3.9, and the last from Lemma 3.7. □

Thus, Theorem A follows since having finite asymptotic dimension implies the existence of a coarse embedding into a Hilbert space [34; 65].

Remark 3.12 Similarly, we can show that $\mathcal{D}: \mathcal{N}^{\leq n} \rightarrow [\mathbb{R}]^{n^2}$ is coarsely $(n^2)^{n^2}$ -to-1 surjective map and so $\text{asdim } \mathcal{N}^{\leq n} \leq \text{asdim } [\mathbb{R}]^{\leq n^2} \leq n^2$ by Theorems 3.9 and 3.5. Hence,

$$\frac{1}{2}n(n-1) \leq \text{asdim } \mathcal{N}^{\leq n} \leq n^2.$$

In particular, $\mathcal{N}^{\leq n}$ can be coarsely embedded into a Hilbert space.

4 Coarse nonembeddability into Hilbert spaces

Let us now recall some definitions and results coming from coarse geometry as they will be the crucial stepping stones to prove our main results.

Suppose that the map $\psi: (X, d_X) \rightarrow (Y, d_Y)$ between metric spaces is a coarse embedding. If two maps $\rho_-: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ and $\rho_+: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ are such that $\rho_- \rightarrow \infty$ and (2) is fulfilled, then we call ρ_- and ρ_+ *control functions*, and we say that ψ is a (ρ_-, ρ_+) -coarse embedding.

Let $\{X_k\}_{k \in \mathbb{N}}$ be a sequence of metric spaces and X be another metric space. We say that a family of maps $\{i_k: X_k \rightarrow X\}_{k \in \mathbb{N}}$ is a *coarse embedding* if there exist two maps $\rho_-, \rho_+: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ such that i_k is a (ρ_-, ρ_+) -coarse embedding for every $k \in \mathbb{N}$. We say that X contains a coarse disjoint union of $\{X_k\}_{k \in \mathbb{N}}$ if there exists a coarse embedding $\{i_k: X_k \rightarrow X\}$ such that

$$\text{dist}(i_n(X_n), i_m(X_m)) \xrightarrow{m+n \rightarrow \infty} \infty.$$

Let us recall that a Banach space $(A, \|\cdot\|)$ is *uniformly convex* if, for every $0 < \varepsilon \leq 2$, there exists $\delta > 0$ so that, for any two vectors $x, y \in A$ with $\|x\| = \|y\| = 1$, the condition $\|x - y\| \geq \varepsilon$ implies that $\|(x + y)/2\| \leq 1 - \delta$. Hilbert spaces are, in particular, uniformly convex Banach spaces.

Theorem 4.1 [39] *There exists a sequence $\{X_k\}_{k \in \mathbb{N}}$ of finite metric spaces such that, if a metric space X contains a coarse disjoint union of $\{X_k\}_{k \in \mathbb{N}}$, then X cannot be coarsely embedded into any uniformly convex Banach space.*

The goal of this section is to prove the following result.

Theorem 4.2 $\mathcal{EH}_1^{<\omega}$ *cannot be coarsely embedded into any uniformly convex Banach space.*

Hence, **Theorem B** immediately follows thanks to **Theorem 2.8** since a composite of coarse embeddings is still a coarse embedding.

To prove **Theorem 4.2**, we intend to apply **Theorem 4.1**. Following the approach used in [63], let us first isometrically embed an arbitrary finite metric space into a more manageable space.

Lemma 4.3 (Kuratowski embedding) *For every finite metric space X , there are $m, n \in \mathbb{N} \setminus \{0\}$ such that X can be isometrically embedded into $([0, m]^n, d_m^n)$, where $d_m^n((x_i)_i, (y_i)_i) = \max_{i=1, \dots, n} |x_i - y_i|$ for every $(x_i)_i, (y_i)_i \in [0, m]^n$.*

Therefore, in order to apply **Theorem 4.1**, we show that $\mathcal{EH}_1^{<\omega}$ contains a coarse disjoint union of the family $\{[0, m]^n \mid m, n \in \mathbb{N} \setminus \{0\}\}$. We intend to define the coarse embeddings $\varphi_m^n: [0, m]^n \rightarrow [\mathbb{R}]^{<\omega}$ recursively. To do it, let us fix a bijection $T: (\mathbb{N} \setminus \{0\})^2 \rightarrow \mathbb{N} \setminus \{0\}$ defined as follows: for every pair $(m, n) \in \mathbb{N}$,

$$T(m, n) = m + \sum_{i=2}^{m+n-1} (i-1) = m + \frac{1}{2}(m+n-2)(m+n-1).$$




Figure 2: A representation of the map $T^{-1}: \mathbb{N} \setminus \{0\} \rightarrow (\mathbb{N} \setminus \{0\})^2$.

We represent in Figure 2 the sequence of the points $T(m, n)$. By construction, for every two pairs $(m, n), (m', n') \in \mathbb{N} \setminus \{0\}$, $m + n \leq m' + n'$ provided that $T(m, n) \leq T(m', n')$. For the sake of simplicity, for two pairs $(m, n), (m', n') \in (\mathbb{N} \setminus \{0\})^2$, let us write $(m, n) \preceq (m', n')$ if $T(m, n) \leq T(m', n')$, and $(m, n) \prec (m', n')$ if $(m, n) \preceq (m', n')$ and $(m, n) \neq (m', n')$.

To construct the maps φ_m^n , we need various parameters. Let us define a sequence $\{a_i(m)\}_{i \in \mathbb{N} \setminus \{0\}}$ of positive real values depending on m as

$$a_i(m) = 4m(i - 1).$$

Furthermore, for every $m, n \in \mathbb{N} \setminus \{0\}$, we inductively construct

$$D(m, n) = \max\{4m(n + 2), \max_{(m', n') \prec (m, n)} D(m', n') + m + 2^{T(m, n)}\}.$$

Thus, $D(1, 1) = 12$ (according to the notation that $\max \emptyset = -\infty$), and $D(m, n) \geq 4m(n + 2) = a_n(m) + 12m$.

Let us define, for every $m, n \in \mathbb{N} \setminus \{0\}$, a map $\varphi_m^n: [0, m]^n \rightarrow [\mathbb{R}]^{=n+1}$ as follows: for every $(x_i)_i \in [0, m]^n$,

$$\varphi_m^n((x_i)_i) = \{a_i(m) + x_i \mid i = 1, \dots, n\} \cup \{D(m, n)\}$$

(see Figure 3). This construction should be compared with that provided in [63]. The crucial difference is the last point $D(m, n)$. Its purpose is to disincentivise the action of isometries and conveniently increase the diameter of the image of φ_m^n (see Lemma 4.4).

Lemma 4.4 *The map $\varphi_m^n: ([0, m]^n, d_m^n) \rightarrow \mathcal{EH}_1^{<\omega}$ is a coarse embedding whose control functions are independent of m and n . More precisely, φ_m^n is a (ρ_-, ρ_+) -coarse embedding, where $\rho_-: x \mapsto x/2$ and $\rho_+ = \text{id}$. Furthermore, for every $(x_i)_i \in [0, m]^n$, we have $\text{diam } \varphi_m^n((x_i)_i) \in [D(m, n) - m, D(m, n)]$.*

Proof It is easy to see that

$$d_{\text{EH}}(\varphi_m^n((x_i)_i), \varphi_m^n((y_i)_i)) \leq d_{\text{H}}(\varphi_m^n((x_i)_i), \varphi_m^n((y_i)_i)) = d_m^n((x_i)_i, (y_i)_i)$$




Figure 3: A representation of the images of the images of two points $(x_i)_i, (y_i)_i \in [0, m]^n$ along φ_m^n . The subset $\varphi_m^n((x_i)_i)$ is given by the red dots and the black dot, while $\varphi_m^n((y_i)_i)$ consists of the blue dots and the black dot. In the picture, we can see that $d_H(\varphi_m^n((x_i)_i), \varphi_m^n((y_i)_i)) = d_m^n((x_i)_i, (y_i)_i)$.

for every $(x_i)_i, (y_i)_i \in [0, m]^n$ (see also Figure 3). We want to prove that ρ_- is the other control function. Let $f \in \text{Isom}(\mathbb{R}^d)$ be an isometry such that

$$(3) \quad d_H(\varphi_m^n((x_i)_i), f(\varphi_m^n((y_i)_i))) \leq d_m^n((x_i)_i, (y_i)_i) \leq m.$$

The isometry f is the composite of a translation g and a rotation h . If $n = 1$, then $|\varphi_m^n((x_i)_i)| = 2$, and so, without loss of generality, h can be taken as the identity.

Claim 4.5 Assume that $n \geq 2$. Then, $h = \text{id}$.

Proof Suppose, by contradiction, that $h \neq \text{id}$. According to (3), and because of the definition of φ_m^n ,

$$(4) \quad |f(D(m, n)) - (a_1(m) + x_1)| \leq m.$$

Since $|(a_1(m) + x_1) - (a_2(m) + x_1)| \geq |a_1(m) - a_2(m)| - m > 2m$, (4) implies that

$$|f(D(m, n)) - (a_2(m) + x_2)| > m.$$

Therefore, because of (3),

$$(5) \quad |(a_2(m) + x_2) - f(a_n(m) + y_n)| \leq m.$$

Using the fact that f is an isometry, the triangular inequality, (4) and (5), the following chain of inequalities descends:

$$\begin{aligned} 11m &\leq |D(m, n) - a_n(m)| - m \leq |D(m, n) - (a_n(m) + y_n)| = |f(D(m, n)) - f(a_n(m) + y_n)| \\ &\leq |f(D(m, n)) - (a_1(m) + x_1)| + |(a_1(m) + x_1) - (a_2(m) + x_2)| + |(a_2(m) + x_2) - f(a_n(m) + y_n)| \\ &\leq m + 5m + m = 7m. \end{aligned}$$

Thus, we obtain a contradiction. \square

We can now assume that the map $f = g$ is a translation. Using the triangular inequality, and since $a_{i+1}(m) - a_i(m) = 4m$ for every $i \in \{1, \dots, n\}$, it can be easily check that

$$|(a_i(m) + x_i) - f(a_i(m) + y_i)| \leq m \quad \text{for every } i \in \{1, \dots, n\} \quad \text{and} \quad |D(m, n) - f(D(m, n))| \leq m.$$

The point $D(m, n)$, common to both $\varphi_m^n((x_i)_i)$ and $\varphi_m^n((y_i)_i)$, misaligns as soon as f is nontrivial. Therefore, f creates a trade-off between the misalignment just described and a better alignment of the other n pairs of points $(a_i(m) + x_i)$ and $a_i(m) + y_i$, for every $i \in \{1, \dots, n\}$. Thus, the best Hausdorff distance that can be achieved is at most $d_H(\varphi_m^n((x_i)_i), \varphi_m^n((y_i)_i))/2 = d_m^n((x_i)_i, (y_i)_i)/2$, and so the claim.

The final assertion trivially follows from the definition of φ_m^n . □

The following result is immediate, but it is an important step in the proof of [Theorem 4.2](#).

Lemma 4.6 *If \mathcal{Y} and \mathcal{Z} are two families of finite subsets of a metric space X ,*

$$\text{dist}_{d_H}(\mathcal{Y}, \mathcal{Z}) \geq \frac{1}{2} \inf_{\substack{Y \in \mathcal{Y} \\ Z \in \mathcal{Z}}} |\text{diam } Y - \text{diam } Z|.$$

Furthermore, if $X = \mathbb{R}^d$, then

$$\text{dist}_{d_{EH}}(\mathcal{Y}, \mathcal{Z}) \geq \frac{1}{2} \inf_{\substack{Y \in \mathcal{Y} \\ Z \in \mathcal{Z}}} |\text{diam } Y - \text{diam } Z|.$$

Proof Since, for every pair of subsets Y and Z of X ,

$$d_H(Y, Z) \geq \frac{1}{2} |\text{diam } Y - \text{diam } Z|,$$

the first inequality is immediate. The second one follows from the fact that an isometry's action does not change the diameter of a subset. □

As a consequence of [Lemma 4.6](#), if \mathcal{Y} and \mathcal{Z} are two families of finite subsets of \mathbb{R}^d with the property that, for every $Y \in \mathcal{Y}$ and $Z \in \mathcal{Z}$, $\text{diam } Y \geq \text{diam } Z$,

$$(6) \quad \text{dist}_{d_{EH}}(\mathcal{Y}, \mathcal{Z}) \geq \frac{1}{2} (\inf_{Y \in \mathcal{Y}} \text{diam } Y - \sup_{Z \in \mathcal{Z}} \text{diam } Z).$$

We now have the tools to show the main result of the section, [Theorem 4.2](#), and its consequence, [Theorem B](#).

Proof of Theorem 4.2 We intend to use Lafforgue's result ([Theorem 4.1](#)), and, thanks to [Lemma 4.3](#), we need to show that $\mathcal{EH}_1^{<\omega}$ contains a coarse disjoint union of $\{([0, m]^n, d_m^n)\}_{m, n \in \mathbb{N} \setminus \{0\}}$. According to [Lemma 4.4](#), $\{\varphi_m^n : [0, m]^n \rightarrow \mathcal{EH}_1^{<\omega}\}$ is a coarse embedding. It remains to show that

$$\text{dist}_{d_{EH}}(\varphi_m^n([0, m]^n), \varphi_{m'}^{n'}([0, m']^{n'})) \xrightarrow{T(m, n) + T(m', n') \rightarrow \infty} \infty.$$

Without loss of generality, we assume that $(m', n') \prec (m, n)$. Then, according to [Lemma 4.4](#), (6), and the definition of $D(m, n)$,

$$(7) \quad \text{dist}_{d_{EH}}(\varphi_m^n([0, m]^n), \varphi_{m'}^{n'}([0, m']^{n'})) \geq \frac{1}{2} (D(m, n) - D(m', n') - m) \geq 2^{T(m, n) - 1}.$$

From (7), the desired result descends. □

5 Bi-Lipschitz nonembeddability into finite-dimensional spaces

Let us consider the isometry classes of all finite subsets of the real line with diameter at most R . This set can be identified with the isometry classes of elements in $[[0, R]]^{<\omega}$. Let us denote by $\mathcal{EH}_{[0, R]}^{<\omega}$ this space equipped with the Euclidean–Hausdorff distance.

Since, for every pair of metric spaces $X, Y \in \mathcal{GH}_{\leq R}^{<\omega}$,

$$d_{\text{GH}}(X, Y) \leq \frac{1}{2} \max\{\text{diam } X, \text{diam } Y\} \leq \frac{1}{2} R,$$

the diameter of $\mathcal{GH}_{\leq R}^{<\omega}$ is finite. Therefore, it can be trivially coarsely embedded into a one-point metric space. However, we can still provide nonembeddability results if we restrict the class of embeddings. More precisely, we prove the following.

Theorem 5.1 $\mathcal{EH}_{[0, R]}^{<\omega}$ cannot be bi-Lipschitz embedded into a finite-dimensional Hilbert space.

Immediately, we can deduce [Theorem C](#) since [Theorem 2.8](#) provides a bi-Lipschitz embedding.

To prove [Theorem 5.1](#), we use the Assouad dimension.

Definition 5.2 Given a metric space (X, d) , a subset $E \subseteq X$ and $r > 0$, we denote by $N_r(E)$ the least number of open balls of radius less or equal to r that cover E . Then, the *Assouad dimension* of X [\[5; 10\]](#) is

$$\dim_A X = \inf\{\alpha > 0 \mid \exists C > 0 : \forall r > 0, \forall \beta \in (0, 1], \sup_{x \in X} N_{\beta r}(B_d^o(x, r)) < C\beta^{-\alpha}\},$$

where $B_d^o(x, r)$ denotes the open ball centred in x with radius r .

This dimension notion was introduced precisely to prove obstructions to bi-Lipschitz embed metric spaces into an Euclidean space. In particular, the following properties lead to the desired conclusion (see, for example, [\[57\]](#)):

- If $\varphi: X \rightarrow Y$ is a bi-Lipschitz embedding between metric spaces, $\dim_A X = \dim_A \text{im}(\varphi) \leq \dim_A Y$.
- For every $n \in \mathbb{N}$, $\dim_A \mathbb{R}^n = n$.

Therefore, once we prove that $\dim_A \mathcal{EH}_{[0, R]}^{<\omega} = \infty$ ([Proposition 5.3](#)), [Theorem 5.1](#) immediately follows.

Let us mention that the same strategy was used in [\[16\]](#) to prove that spaces of persistence diagrams cannot be bi-Lipschitz embedded into a finite-dimensional Hilbert space.

Proposition 5.3 For every $R > 0$, $\dim_A \mathcal{EH}_{[0, R]}^{<\omega} = \infty$.

Proof We want to provide, for every $\alpha > 0$ and every $C > 0$, a radius $r > 0$, a constant $\beta \in (0, 1]$, a finite subset A of $[0, R]$ and $M = \lceil C\beta^{-\alpha} + 1 \rceil$ -many finite subsets A_1, \dots, A_M of $[0, R]$ with the following properties: $d_{\text{EH}}(A, A_i) < r$ and $d_{\text{EH}}(A_i, A_j) > 2\beta r$ for every $i, j \in \{1, \dots, M\}$. Therefore, the open ball centred in A with radius r cannot be covered by fewer than M -many open balls with radius βr (A_i and A_j are contained in the same ball only if $i = j$), which will conclude the proof since $M > C\beta^{-\alpha}$.




Figure 4: A representation of the subsets A , A_1 and A_i defined in the proof of Proposition 5.3. The distinctive points $l - s \in A_1$ and $il - s \in A_i$ are emphasised in red. The light grey strips are meant to visualise the fact that $d_H(A, A_1) < r$ and $d_H(A, A_i) < r$.

Define $l = \frac{R}{2(M+1)}$, $r = \frac{l}{6}$, $s = \frac{2r}{3}$ and $\beta = \frac{1}{2}$. We construct the subsets A and A_i as

$$\begin{aligned} A &= \{j \cdot l \mid j \in \{0, \dots, M+1\}\} \cup \{R\}, \\ A_i &= \{j \cdot l + s \mid j \in \{1, \dots, M\} \setminus \{i\}\} \cup \{i \cdot l - s\} \cup \{0, \frac{1}{2}R, R\}. \end{aligned}$$

The subset A and A_i are represented in Figure 4. It is clear that $d_{\text{EH}}(A, A_i) < r$ since $s < r$ (actually, $d_H(A, A_i) = s < r$). It remains to show that, for every pair of distinct indices $i, j \in \{1, \dots, M\}$, $d_{\text{EH}}(A_i, A_j) \geq 2s > r$.

Let $i, j \in \{1, \dots, M\}$ be two distinct indices. Assume, by contradiction, that $d_{\text{EH}}(A_i, A_j) < 2s$, and let f be an isometry of \mathbb{R} such that $d_H(A_i, f(A_j)) < 2s$. Adapting the argument of Claim 4.5, we can assume that f is a translation.

For the sake of simplicity, for every $k \in \{1, \dots, n\}$, we name the points of A_k as

$$\begin{aligned} a_0^k &= 0, & a_1^k &= l + s, & \dots, & a_{k-1}^k &= (k-1)l + s, & a_k^k &= kl - s, \\ a_{k+1}^k &= (k+1)l + s, & \dots, & a_M^k &= Ml + s, & a_{M+1}^k &= \frac{1}{2}R, & a_{M+2}^k &= R. \end{aligned}$$

Following the strategy used in the proof of Lemma 4.4, we can show that, for every $k \in \{0, \dots, M+2\}$,

$$(8) \quad |a_k^i - f(a_k^j)| < 2s.$$

Assume, without loss of generality, that $i < j$. Then, in particular,

$$(9) \quad |a_i^j - a_j^j| = (j-i)l - 2s \quad \text{and} \quad |a_i^i - a_j^i| = (j-i)l + 2s.$$

Using the triangular inequality, (8), (9) and the fact that f is an isometry we obtain

$$\begin{aligned} (j-i)l + 2s &= |a_i^i - a_j^i| \leq |a_i^i - f(a_i^j)| + |f(a_i^j) - f(a_j^j)| + |f(a_j^j) - a_j^i| \\ &< |a_i^j - a_j^j| + 4s = (j-i)l + 2s. \end{aligned}$$

Hence, we have found a contradiction. \square

Remark 5.4 As a byproduct of the proof of [Proposition 5.3](#), we obtain that the Assouad dimension of the family of all finite subsets of an interval equipped with the Hausdorff distance is infinite. Indeed, in the notation of the mentioned proof, $d_H(A, A_i) = s < r$, but $d_H(A_i, A_j) = 2s > r$ for every pair of distinct indices $i, j \in \{1, \dots, M\}$.

5.1 Questions about bi-Lipschitz embeddability and Assouad dimension

Let us conclude the presentation with a discussion about potential future research directions concerning the bi-Lipschitz embeddability of the Gromov–Hausdorff space. First of all, [Theorem C](#) leaves the following question open.

Question 5.5 *Can $\mathcal{GH}_{\leq R}^{<\omega}$ be bi-Lipschitz embedded into an infinite-dimensional Hilbert space?*

Furthermore, it is natural to ask what the embeddability properties are if we bound the cardinality of the metric spaces as in [Section 3](#). Let us define $\mathcal{GH}_{\leq R}^{\leq n} = \mathcal{GH}^{\leq n} \cap \mathcal{GH}_{\leq R}^{<\omega}$ for $n \in \mathbb{N}$ and $R > 0$.

Question 5.6 *Can $\mathcal{GH}^{\leq n}$ and $\mathcal{GH}_{\leq R}^{\leq n}$ be bi-Lipschitz embedded into a (finite-dimensional) Hilbert space?*

To approach [Question 5.6](#), we may investigate the Assouad dimension of those spaces. The inequalities

$$\dim_A \mathcal{GH}^{\leq n} \geq \dim_A \mathcal{GH}_{\leq R}^{\leq n} \geq \frac{1}{2}n(n-1)$$

can be derived similarly to the proof of [Lemma 3.4](#). Indeed, according to [\[35, Theorem 4.1\]](#), $\mathcal{GH}_{\leq R}^{\leq n}$ contains a subspace isometric to an open ball in $\mathbb{R}^{n(n-1)/2}$ of sufficiently small radius, which has Assouad dimension $n(n-1)/2$ [\[57, Lemma 9.6\(iii\)\]](#). Hence, monotonicity and bi-Lipschitz invariance of the dimension imply the desired estimate. However, to the best of the author’s knowledge, no upper bounds are known for the dimension of those spaces, and the following questions remain open.

Question 5.7 *What are $\dim_A \mathcal{GH}^{\leq n}$ and $\dim_A \mathcal{GH}_{\leq R}^{\leq n}$? Are they finite?*

A connection between [Questions 5.7](#) and [5.6](#) has already been exploited to deduce [Theorem C](#), namely, infinite Assouad dimension prevents the existence of bi-Lipschitz embeddings into finite-dimensional Hilbert spaces. However, unlike the situation described for asymptotic dimension and coarse embeddings, a positive answer to the second part of [Question 5.7](#) does not imply the existence of a bi-Lipschitz embedding into some \mathbb{R}^n . Indeed, having finite Assouad dimension is not a sufficient condition for the existence of a bi-Lipschitz embedding even into some infinite-dimensional Hilbert space [\[38; 40; 53; 60\]](#). However, some positive results can be proved at the cost of modifying the original metric space. For a metric space (X, d) and $0 < \varepsilon < 1$, the ε -snowflaking of X is the metric space (X, d^ε) , where $d^\varepsilon(x, y) = (d(x, y))^\varepsilon$.

Theorem 5.8 (Assouad embedding theorem [\[5\]](#)) *If (X, d) is a metric space with finite Assouad dimension, then, for every $0 < \varepsilon < 1$, there exists a bi-Lipschitz embedding of (X, d^ε) into \mathbb{R}^n for some n depending only on $\dim_A X$ and ε .*

Let us also mention that, if $\frac{1}{2} < \varepsilon < 1$, the parameter n in the Assouad embedding theorem can be chosen independently from ε ([51]; see also [21] for an explicit map construction). We address the interested reader to the monographs [28; 57] for more details. A positive answer to the second part of Question 5.7 could then motivate the search for computable bi-Lipschitz embeddings of the ε -snowflaking of $\mathcal{GH}^{\leq n}$ or $\mathcal{GH}_{\leq R}^{\leq n}$ into some finite-dimensional Hilbert space, further tightening the connection between computational topology and dimension theory.

References

- [1] H Adams, J Bush, N Clause, F Frick, M Gómez, M Harrison, R A Jeffs, E Lagoda, S Lim, F Mémoli, M Moy, N Sadovek, M Superdock, D Vargas, Q Wang, L Zhou, *Gromov–Hausdorff distances, Borsuk–Ulam theorems, and Vietoris–Rips complexes*, preprint (2023) [arXiv 2301.00246](https://arxiv.org/abs/2301.00246)
- [2] P K Agarwal, K Fox, A Nath, A Sidirooulos, Y Wang, *Computing the Gromov–Hausdorff distance for metric trees*, ACM Trans. Algorithms 14 (2018) art. id. 24 [MR](#)
- [3] S A Antonyan, *The Gromov–Hausdorff hyperspace of a Euclidean space*, Adv. Math. 363 (2020) art. id. 106977 [MR](#)
- [4] S A Antonyan, *The Gromov–Hausdorff hyperspace of a Euclidean space, II*, Adv. Math. 393 (2021) art. id. 108055 [MR](#)
- [5] P Assouad, *Plongements lipschitziens dans \mathbb{R}^n* , Bull. Soc. Math. France 111 (1983) 429–448 [MR](#)
- [6] D Bate, A L Garcia Pulido, *Bi-Lipschitz embeddings of the space of unordered m -tuples with a partial transportation metric*, Math. Ann. 390 (2024) 3109–3131 [MR](#)
- [7] G Bell, A Dranishnikov, *Asymptotic dimension*, Topology Appl. 155 (2008) 1265–1296 [MR](#)
- [8] G S Bloom, *A counterexample to a theorem of S Picard*, J. Combinatorial Theory Ser. A 22 (1977) 378–379 [MR](#)
- [9] L M Blumenthal, *Theory and applications of distance geometry*, Clarendon, Oxford (1953) [MR](#)
- [10] M G Bouligand, *Ensembles impropre et nombre dimensionnel*, Bull. Sci. Math. 52 (1928) 320–344, 361–376
- [11] P Bubenik, A Wagner, *Embeddings of persistence diagrams into Hilbert spaces*, J. Appl. Comput. Topol. 4 (2020) 339–351 [MR](#)
- [12] D Burago, Y Burago, S Ivanov, *A course in metric geometry*, Graduate Studies in Math. 33, Amer. Math. Soc., Providence, RI (2001) [MR](#)
- [13] G Carlsson, *Topology and data*, Bull. Amer. Math. Soc. 46 (2009) 255–308 [MR](#)
- [14] G Carlsson, F Mémoli, *Persistent clustering and a theorem of J Kleinberg*, preprint (2008) [arXiv 0808.2241](https://arxiv.org/abs/0808.2241)
- [15] G Carlsson, F Mémoli, *Characterization, stability and convergence of hierarchical clustering methods*, J. Mach. Learn. Res. 11 (2010) 1425–1470 [MR](#)
- [16] M Carrière, U Bauer, *On the metric distortion of embedding persistence diagrams into separable Hilbert spaces*, from “35th International Symposium on Computational Geometry” (G Barequet, Y Wang, editors), Leibniz Int. Proc. Inform. 129, Schloss Dagstuhl, Wadern (2019) art. id. 21 [MR](#)

- [17] **F Chazal, D Cohen-Steiner, L J Guibas, F Mémoli, S Y Oudot**, *Gromov–Hausdorff stable signatures for shapes using persistence*, Computer Graphics Forum 28 (2009) 1393–1403
- [18] **F Chazal, V de Silva, S Oudot**, *Persistence stability for geometric complexes*, Geom. Dedicata 173 (2014) 193–214 [MR](#)
- [19] **S Chowdhury, F Mémoli**, *A functorial Dowker theorem and persistent homology of asymmetric networks*, J. Appl. Comput. Topol. 2 (2018) 115–175 [MR](#)
- [20] **S Chowdhury, F Mémoli**, *Distances and isomorphism between networks: stability and convergence of network invariants*, J. Appl. Comput. Topol. 7 (2023) 243–361 [MR](#)
- [21] **G David, M Snipes**, *A constructive proof of the Assouad embedding theorem with bounds on the dimension*, preprint (2012) [arXiv 1211.3223](#)
- [22] **A N Dranishnikov, G Gong, V Lafforgue, G Yu**, *Uniform embeddings into Hilbert space and a question of Gromov*, Canad. Math. Bull. 45 (2002) 60–70 [MR](#)
- [23] **J Dydak, Ž Virk**, *Preserving coarse properties*, Rev. Mat. Complut. 29 (2016) 191–206 [MR](#)
- [24] **J Dydak, T Weighill**, *Monotone-light factorizations in coarse geometry*, Topology Appl. 239 (2018) 160–180 [MR](#)
- [25] **H Edelsbrunner, J L Harer**, *Computational topology: an introduction*, Amer. Math. Soc., Providence, RI (2010) [MR](#)
- [26] **H Edelsbrunner, T Heiss, V Kurlin, P Smith, M Wintraecken**, *The density fingerprint of a periodic point set*, from “37th International Symposium on Computational Geometry” (K Buchin, E Colin de Verdière, editors), Leibniz Int. Proc. Inform. 189, Schloss Dagstuhl, Wadern (2021) art. id. 32 [MR](#)
- [27] **D A Edwards**, *The structure of superspace*, from “Studies in topology” (N M Stavrakas, K R Allen, editors), Academic, New York (1975) 121–133 [MR](#)
- [28] **J M Fraser**, *Assouad dimension and fractal geometry*, Cambridge Tracts in Mathematics 222, Cambridge Univ. Press (2021) [MR](#)
- [29] **A Garber, Ž Virk, N Zava**, *On the metric spaces of lattices and periodic point sets*, preprint (2023) [arXiv 2310.07594](#)
- [30] **B Giunti, J Lazovskis, B Rieck**, *DONUT: database of original and non-theoretical uses of topology* (2022) Available at <https://donut.topology.rocks>
- [31] **M Gromov**, *Structures métriques pour les variétés riemanniennes*, Textes Mathématiques 1, CEDIC, Paris (1981) [MR](#)
- [32] **M Gromov**, *Asymptotic invariants of infinite groups*, from “Geometric group theory, II” (G A Niblo, M A Roller, editors), London Math. Soc. Lecture Note Ser. 182, Cambridge Univ. Press (1993) 1–295 [MR](#)
- [33] **K Hess**, *Topological adventures in neuroscience*, from “Topological data analysis—the Abel Symposium 2018” (N A Baas, G E Carlsson, G Quick, M Szymik, M Thaule, editors), Abel Symp. 15, Springer (2020) 277–305 [MR](#)
- [34] **N Higson, J Roe**, *Amenable group actions and the Novikov conjecture*, J. Reine Angew. Math. 519 (2000) 143–153 [MR](#)
- [35] **S Iliadis, A O Ivanov, A A Tuzhilin**, *Local structure of Gromov–Hausdorff space, and isometric embeddings of finite metric spaces into this space*, Topology Appl. 221 (2017) 393–398 [MR](#)
- [36] **N J Kalton, M I Ostrovska**, *Distances between Banach spaces*, Forum Math. 11 (1999) 17–48 [MR](#)

- [37] **J Kucab, M Zarichnyi**, *On asymptotic power dimension*, Topology Appl. 201 (2016) 124–130 [MR](#)
- [38] **T J Laakso**, *Plane with A_∞ -weighted metric not bi-Lipschitz embeddable to \mathbb{R}^N* , Bull. London Math. Soc. 34 (2002) 667–676 [MR](#)
- [39] **V Lafforgue**, *Un renforcement de la propriété (T)*, Duke Math. J. 143 (2008) 559–602 [MR](#)
- [40] **U Lang, C Plaut**, *Bilipschitz embeddings of metric spaces into space forms*, Geom. Dedicata 87 (2001) 285–307 [MR](#)
- [41] **S Lim, F Mémoli, Z Smith**, *The Gromov–Hausdorff distance between spheres*, Geom. Topol. 27 (2023) 3733–3800 [MR](#)
- [42] **S Majhi, J Vitter, C Wenk**, *Approximating Gromov–Hausdorff distance in Euclidean space*, Comput. Geom. 116 (2024) art. id. 102034 [MR](#)
- [43] **F Mémoli**, *On the use of Gromov–Hausdorff distances for shape comparison*, from “Eurographics Symposium on Point-Based Graphics” (M Botsch, R Pajarola, B Chen, M Zwicker, editors), The Eurographics Association (2007) 81–90
- [44] **F Mémoli**, *Gromov–Hausdorff distances in Euclidean spaces*, from “2008 IEEE Computer Society Conference on Computer Vision and Pattern Recognition Workshops” (2008) 1–8
- [45] **F Mémoli**, *Some properties of Gromov–Hausdorff distances*, Discrete Comput. Geom. 48 (2012) 416–440 [MR](#)
- [46] **F Mémoli, G Sapiro**, *Comparing point clouds*, from “SGP ’04: Proceedings of the 2004 Eurographics/ACM SIGGRAPH Symposium on Geometry Processing”, ACM, New York (2004) 32–40
- [47] **F Mémoli, G Sapiro**, *A theoretical and computational framework for isometry invariant recognition of point cloud data*, Found. Comput. Math. 5 (2005) 313–347 [MR](#)
- [48] **A Mitra, Ž Virk**, *The space of persistence diagrams on n points coarsely embeds into Hilbert space*, Proc. Amer. Math. Soc. 149 (2021) 2693–2703 [MR](#)
- [49] **A Mitra, Ž Virk**, *Geometric embeddings of spaces of persistence diagrams with explicit distortions*, preprint (2024) [arXiv 2401.05298](#)
- [50] **T Miyata, Ž Virk**, *Dimension-raising maps in a large scale*, Fund. Math. 223 (2013) 83–97 [MR](#)
- [51] **A Naor, O Neiman**, *Assouad’s theorem with dimension independent of the snowflaking*, Rev. Mat. Iberoam. 28 (2012) 1123–1142 [MR](#)
- [52] **P W Nowak, G Yu**, *Large scale geometry*, European Math. Society, Zürich (2012) [MR](#)
- [53] **P Pansu**, *Métriques de Carnot–Carathéodory et quasiisométries des espaces symétriques de rang un*, Ann. of Math. 129 (1989) 1–60 [MR](#)
- [54] **P Petersen**, *Riemannian geometry*, Graduate Texts in Mathematics 171, Springer (1998) [MR](#)
- [55] **N Pritchard, T Weighill**, *Coarse embeddability of Wasserstein space and the space of persistence diagrams*, Discrete Comput. Geom. 74 (2025) 358–373 [MR](#)
- [56] **T M Radul, O Shukel**, *Functors of finite degree and asymptotic dimension*, Mat. Stud. 31 (2009) 204–206 [MR](#)
- [57] **J C Robinson**, *Dimensions, embeddings, and attractors*, Cambridge Tracts in Mathematics 186, Cambridge Univ. Press (2011) [MR](#)

- [58] **J Roe**, *Lectures on coarse geometry*, University Lecture Series 31, Amer. Math. Soc., Providence, RI (2003) [MR](#)
- [59] **F Schmiedl**, *Computational aspects of the Gromov–Hausdorff distance and its application in non-rigid shape matching*, Discrete Comput. Geom. 57 (2017) 854–880 [MR](#)
- [60] **S Semmes**, *On the nonexistence of bi-Lipschitz parameterizations and geometric problems about A_∞ -weights*, Rev. Mat. Iberoamericana 12 (1996) 337–410 [MR](#)
- [61] **A A Tuzhilin**, *Lectures on Hausdorff and Gromov–Hausdorff distance geometry*, preprint (2020) [arXiv 2012.00756v1](#)
- [62] **A Wagner**, *Nonembeddability of persistence diagrams with $p > 2$ Wasserstein metric*, Proc. Amer. Math. Soc. 149 (2021) 2673–2677 [MR](#)
- [63] **T Weighill, T Yamauchi, N Zava**, *Coarse infinite-dimensionality of hyperspaces of finite subsets*, Eur. J. Math. 8 (2022) 335–355 [MR](#)
- [64] **G Yu**, *The Novikov conjecture for groups with finite asymptotic dimension*, Ann. of Math. 147 (1998) 325–355 [MR](#)
- [65] **G Yu**, *The coarse Baum–Connes conjecture for spaces which admit a uniform embedding into Hilbert space*, Invent. Math. 139 (2000) 201–240 [MR](#)
- [66] **N Zava**, *Stability of the q -hyperconvex hull of a quasi-metric space*, preprint (2022) [arXiv 2208.10619](#)

Institute of Science and Technology Austria
Klosterneuburg, Austria

nicolo.zava@gmail.com

Received: 30 June 2024 Revised: 4 October 2024

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre etnyre@math.gatech.edu Georgia Institute of Technology	Kathryn Hess kathryn.hess@epfl.ch École Polytechnique Fédérale de Lausanne
--	--

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Markus Land	LMU München markus.land@math.lmu.de
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Norihiro Minami	Yamato University minami.norihiro@yamato-u.ac.jp
Octav Cornea	Université de Montréal cornea@dms.umontreal.ca	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futer	Temple University dfuter@temple.edu	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mheden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2025 is US \$760/year for the electronic version, and \$1110/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY
 **mathematical sciences publishers**

nonprofit scientific publishing

<https://msp.org/>

© 2025 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 25

Issue 8 (pages 4437–5174)

2025

Hierarchies for relatively hyperbolic virtually special groups	4437
EDUARD EINSTEIN	
Intersection norms on surfaces and Birkhoff sections for geodesic flows	4499
MARCOS COSSARINI and PIERRE DEHORNOY	
Thin knots and the cabling conjecture	4547
ROBERT DEYESO III	
Linear linkless embeddings: proof of a conjecture by Sachs	4585
LYNN STANFIELD	
Constructing rational homology 3-spheres that bound rational homology 4-balls	4599
LISA LOKTEVA	
Homological stability for the ribbon Higman–Thompson groups	4633
RACHEL SKIPPER and XIAOLEI WU	
A group-theoretic framework for low-dimensional topology, or: how not to study low-dimensional topology?	4667
SARAH BLACKWELL, ROBION KIRBY, MICHAEL KLUG, VINCENT LONGO and BENJAMIN RUPPIK	
Classification of genus-two surfaces in S^3	4719
FILIPPO BARONI	
Meromorphic projective structures: signed spaces, grafting and monodromy	4787
SPANDAN GHOSH and SUBHOJOY GUPTA	
A new twist on modular links from an old perspective	4827
KHANH LE	
Flat fully augmented links are determined by their complements	4839
CHRISTIAN MILICHAP and ROLLAND TRAPP	
BNSR-invariants of surface Houghton groups	4897
NOAH TORGERSON and JEREMY WEST	
Topological symmetry groups of the generalized Petersen graphs	4921
ANGELYNN ÁLVAREZ, ERICA FLAPAN, MARK HUNNELL, JOHN HUTCHENS, EMILLE LAWRENCE, PAUL LEWIS, CANDICE PRICE and RUTH VANDERPOOL	
Crushing surfaces of positive genus	4949
BENJAMIN A BURTON, THIAGO DE PAIVA, ALEXANDER HE and CONNIE ON YU HUI	
Annular links from Thompson's group T	5013
LOUISA LILES	
Realizing pairs of multicurves as cylinders on translation surfaces	5031
JULIET AYGUN, JANET BARKDOLL, AARON CALDERON, JENAVIE LORMAN and THEODORE SANDSTROM	
Involutive Khovanov homology and equivariant knots	5059
TAKETO SANO	
A diagrammatic computation of abelian link invariants	5113
DAVID CIMASONI, LIVIO FERRETTI and JESSICA LIU	
Equivariant double-slice genus, stabilization, and equivariant stabilization	5137
MALCOLM GABBARD	
Coarse and bi-Lipschitz embeddability of subspaces of the Gromov–Hausdorff space into Hilbert spaces	5153
NICOLÒ ZAVA	